

# ISOLATION GUIDANCE FOR GENERAL PUBLIC

**START**

**HAVE YOU TESTED POSITIVE  
FOR COVID-19?**

**YES**

**NO**

**ARE YOU EXPERIENCING  
ANY SYMPTOMS?**

**YES**

**NO**

5 day isolation. (Day 0 is the day you started experiencing symptoms or tested positive, whichever occurred first.) If you have a fever, continue to stay home until your fever resolves for at least 24 hours. Continue to wear a mask around others for additional 5 days.

5 day isolation and then you must wear a mask around others for an additional 5 days.

**YES**

**NO**

**HAVE OR ARE YOU  
EXPERIENCING ANY  
SYMPTOMS?**

**WERE YOU A CLOSE  
CONTACT TO A POSITIVE  
CASE OF COVID-19?**

**YES**

**NO**

You may have something else, like the flu or other infection. Contact your primary care provider. Stay home until feeling better.

Follow quarantine guidance

You can go about normal activities, but continue to practice safety measures.

# QUARANTINE GUIDANCE FOR GENERAL PUBLIC

**START**

**WERE YOU WITHIN 6 FT. OR LESS OF A POSITIVE CASE FOR MORE THAN 15 MINUTES, WITHIN 48 HOURS PRIOR TO THEIR SYMPTOMS STARTING OR TESTING POSITIVE?**

You are not considered a close contact and can resume normal activities.  
If symptoms occur, get tested.

**NO**

**YES**

**ARE YOU VACCINATED?**

**DID YOU RECEIVE A BOOSTER SHOT?**

**YES**

**NO**

**YES OR N/A\***

You do not have to quarantine.  
Wear mask for 10 days following exposure. Best practice would include being tested 5 days after exposure or at first sign of any symptoms.

**NO**

Quarantine for 5 days. Day 0 is the day in which you were around positive case. You must wear a mask around anyone for the next 5 days. If symptoms occur, get tested.

**HAS IT BEEN MORE THAN 6 MONTHS SINCE YOUR LAST PFIZER OR MODERNA VACCINE OR MORE THAN 2 MONTHS SINCE YOUR LAST J&J?**

**YES**

Quarantine for 5 days. Day 0 is the day in which you were around positive case. You must wear a mask around anyone for the next 5 days. If symptoms occur, get tested.

**NO**

You do not have to quarantine.  
Wear mask for 10 days following exposure. Best practice would include being tested 5 days after exposure or at first sign of any symptoms.

\*\* N/A - see CDC guidelines on booster eligibility based on vaccine & age